

Measurement of Change of ^7Be decay rate in Be and Au

Zhiyi LIU, Chengbo LI, Siguang WANG, Jing ZHOU,
Qiuying MENG, Shaojun LU, Shuhua ZHOU

February 8, 2008

Abstract

We have measured the possible change of the decay rate of ^7Be implanted into hosts of natural beryllium and natural gold. No difference between the ^7Be decay rates in the two hosts is observed within the experimental precision of 0.12%. This result implies that change of the decay rate of ^7Be implanted in different materials cannot be simply expected from the electron affinity difference consideration lonely and the lattice structure of the host materials should be taken into account.
PACS:23.40.-s, 71.20.-b

Electron-capture decay rates depend sensitively on the density of atomic electrons within the nucleus. Thus, physically and chemically environmental factors such as pressure, chemical form, magnetic fields, etc. that can alter electron densities, may affect electron-capture decay rates. Since the change of nuclear decay in different environments has fundamental significance, as well as application in nuclear physics, geology and condensed matter physics, such a study is of current interests.^[1, 2, 3, 4, 5] ^7Be is the lightest radioactive nucleus that decays by electron capture with a half-life of ~ 53 days, thereby, it is a good candidate for studying perturbation of nuclear decay rates. Furthermore, the study of decay rate changes for ^7Be has particular significance to the solar neutrino problem, where there is a large discrepancy between theoretical predictions and experimental determinations of the solar neutrino flux.^[3, 6, 7, 8]

The ^7Be decay rate sensitively depends on chemical environment at the nucleus. Changes of decay rates of ^7Be nuclei in different ^7Be compounds have been measured at normal pressure and a maximum change is about 1.5%.^[9] Moreover, this decay rate is also sensitive to physical environments, such as high pressure and host materials in which the ^7Be nucleus is located. Liu et al. have measured a large increase, up to 1%, of the decay rate of ^7Be under a high pressure of 400 kilobars.^[10] Recently Norman et al. measured the decay rate of ^7Be implanted into hosts of lithium fluoride, gold, graphite, boron nitride, tantalum, lithium, and so on. It has been found that the ^7Be decay rate varies by as much as 0.72% from one host to another.^[11, 12, 13, 14] Qualitatively, it has been reported that if a ^7Be atom is implanted in a medium having high electron affinity (EA), as a result of its interaction with nearby atoms of such a medium, the ^7Be atom

would lose a significant fraction of its $2s$ electrons.^[12] The decay rate of ^7Be in a high-EA medium is thereby smaller than that in a low-EA medium. Thus, we would expect that ^7Be implanted in natural gold ($\text{EA}_{\text{Au}}=2.308\text{eV}$ ^[15]) should decay slower up to 0.7% than that in natural beryllium ($\text{EA}_{\text{Be}}=0.19\text{eV}$ ^[15]) according to the result of Ray et al., in which difference up to 0.72% between the decay rates of ^7Be implanted in Au and in Al_2O_3 was observed.^[12]

In this Letter, we report our measurement of possible change of the decay rate of ^7Be implanted into metal foils of natural beryllium and natural gold. The ^7Be nuclei are produced by bombarding a $500\text{ }\mu\text{g}/\text{cm}^2$ -thick foil of lithium fluoride (LiF) with a 3.2 MeV proton beam, with average current $5\text{ }\mu\text{A}$, from the 5SDH-2 tandem accelerator at CNNC Radiation Metrology and Measurement Center, China Institute of Atomic Energy. Thus, ^7Be nuclei produced by the reaction $^7\text{Li}(\text{p}, \text{n})^7\text{Be}$ with recoil energy 1.0 MeV in the forward direction are implanted into beryllium and gold foils placed immediately behind the LiF target. The implantation for each foil lasts about 15 h. As a result of such implantations, ^7Be atoms are expected to be randomly located in the interstitial lattice space of the host media beryllium and gold.

By electron capture the nucleus ^7Be decays to the $3/2^-$ ground state of ^7Li directly with a branching ratio of 89.5% and to the first excited state with a ratio of 10.5%, which decays subsequently to its ground state by emitting a 478-keV gamma ray.^[16] In this experiment, two high-purity coaxial germanium (HP Ge) detectors are used to measure the 478-keV gamma-ray photons. The ^7Be implanted beryllium and gold foils are separately mounted about 2 cm away from the endcaps of two HP Ge detectors. The two detectors, each is surrounded by a graded shield of organic glass-Al-Cu-Pb, are located four meters away from each other to avoid cross detection of the gamma-ray photons from the two sources. The gamma-ray spectra are accumulated with a KODAQ data acquisition system.

In order to decrease as much effect originated from the decays of short-lived isotopes produced in bombarding as possible, we wait for about 40 days to start to acquire experimental data. The two gamma-ray spectra are acquired for successive one-hour intervals and recorded on a computer hard disk. Then this is followed by next interval automatically. After such a measurement has lasted about 70 days, we exchange the positions of the two sources, while keeping other measurement conditions unchanged, and we measure for another 50 days. We extract from each time bin the net peak area of the 478 keV ^7Be gamma-ray and its time information after dead-time correction and time calibrations with the time provided by Hongkong observatory. Figure 1 shows a typical gamma-ray spectrum in a one-hour counting period from the gold sample. The ratio method, as to be described below, is employed to draw the possible difference in the decay rate of ^7Be nuclei implanted in beryllium and gold. Let A_{Be} and A_{Au} denote the numbers of 478 keV gamma-ray photons measured by the corresponding detectors within an interval of t to $(t+1)$ days. Similarly, let A_{Be}^0 and A_{Au}^0 denote the numbers of 478 keV gamma-ray photons measured by the corresponding detectors within an interval of 0 to $t=1$ days. Let λ_{Be} and λ_{Au} denote the decay rate of the ^7Be nuclei in

the beryllium and gold samples respectively and let $\lambda_{Be} = \lambda_{Au} + \Delta\lambda$. Then

$$\ln R(t) = \ln R_0 - \Delta\lambda t \quad (1)$$

where $R(t) = \frac{A_{Be}}{A_{Au}}$, $R_0 = \frac{A_{Be}^0}{A_{Au}^0}$, and $\Delta\lambda$ is the difference in the decay rate.




Figure 1: A typical gamma-ray spectrum obtained in one-hour counting interval from ${}^7\text{Be}$ in the gold sample




Figure 2: Characteristics of $\ln R$ versus time (a) before and (b) after exchanging sources positions. The solid lines are the linear fit of the experimental data.

Figures 2(a) and 2(b) show the experimental data of $\ln R(t)$ versus t and corresponding linear fit lines before and after exchanging the source positions, respectively. The $\Delta\lambda$ value is determined by the slope in each case. The value of $\Delta\lambda$ derived from Figs.

2(a) and 2(b) are $0.3 \times 10^{-5} \pm 0.8 \times 10^{-5}$ and $-0.7 \times 10^{-5} \pm 1.5 \times 10^{-5}$, respectively. By fitting the exponential decrease of A_{Au} versus time t for the two cases, we obtain the weighted value 0.012985 ± 0.000004 for the decay rate of ^7Be implanted in gold and 0.012986 ± 0.000004 in beryllium. Then using λ equal to 0.012986, we obtain the values of $(0.02 \pm 0.06)\%$ before the source exchange and $(-0.05 \pm 0.12)\%$ after the source exchange. This result indicates that within our experimental precision, the large change in the decay rate of ^7Be in Be and Au is not observed and an upper limit 0.12% of this change can be set.

In conclusion, we have performed the first-ever measurement of the possible change in the decay rate of ^7Be implanted in natural beryllium and natural gold. According to the large difference of electron affinity between beryllium and gold, a large change of 0.7% in the decay rate of ^7Be in the two cases is expected. However, within our measurement precision ($\sim \pm 0.06$ and 0.12% for the measurement before and after the source exchange respectively) no difference between the decay rates of ^7Be implanted in beryllium and in gold is observed. This result implies that except electron affinity the lattice structure of the host medium in which the ^7Be atom sits has to be taken into account.

Our experimental result also indicates that the difference between the average electron numbers in L -shell ($2s$ state) of ^7Be atoms in the two samples is smaller than our measurement precision of $\sim 0.1\%$, since the K -shell capture rate should essentially remain unchanged in different environments. According to the calculation of Ray et al., which has taken the lattice structure into account, the average number of $2s$ electrons of ^7Be atoms implanted in natural beryllium sample should be equal to 0.443 while that in the gold sample should be 0.416.^[12, 17, 18] Thus, from Hartree's calculation that the ratio of the square of the beryllium $2s$ electronic state wave function ($2s$ electrons) to that of $1s$ state wave function at the nucleus ($r=0$) is 3.31%,^[19] the decay rate of ^7Be in natural beryllium should be faster than that in gold by 0.045%. However, this value is beyond the reach of our measurement precision.

Acknowledgement

We are grateful to Amlan Ray at Variable Energy Cyclotron Centre of India for useful discussion. We also acknowledge Da-Qing Yuan and Chao-Fan Rong for using their gamma-ray spectroscopy laboratory.

References

- [1] Emery G T 1972 *Ann. Rev. Nucl. Sci.* **22** 165
- [2] Dostal K P, Nagel M and Pabst D 1977 *Z. Naturforsch.* **32a** 345
- [3] Bahcall J N 1996 *Astrophys. J.* **467** 475
- [4] Kerr R A 1999 *Science* **286** 882

- [5] Johlige H W, Aumann D C and Born H J 1970 *Phys. Rev. C* **2** 1616
- [6] Wolfenstein L and Krastev PI 1997 *Phys. Rev. D* **55** 4405
- [7] Chauhan B C and Dev S 1997 *J. Phys. Soc. Jpn.* **66** 917
- [8] Junghans A R et al. 2002 *Phys. Rev. Lett.* **88** 041101
- [9] Huh C-A 1999 *Earth Planet. Sci. Lett.* **171** 325
- [10] Liu L-G and Huh C-A 2000 *Earth Planet. Sci. Lett.* **180** 163
- [11] Jaeger M et al. 1996 *Phys. Rev. C* **54** 423
- [12] Ray A et al. 1999 *Phys. Lett. B* **455** 69
- [13] Norman E B et al. 2001 *Phys. Lett. B* **519** 15
- [14] Souza D J et al. 2002 *J. Nucl. Sci. Technol. Suppl.* **2** (August) 470
- [15] Lide D R (Ed.) 2002 *CRC Handbook of Chemistry and Physics* (Florida and: CRC Press 2002 83rd) vol 83 pp 10-147
- [16] Firestone R B et al. 1996 *Table of Isotopes CD-ROM* (New York and: John Wiley & Sons Inc. 1996 8th.) vol 8 p 272
- [17] Ray A et al. 2002 *Phys. Lett. B* **531** 187
- [18] Ray A et al. 2002 *Phys. Rev. C* **66** 012501(R)
- [19] Hartree D R and Hartree W 1935 *Proc. Roy. Soc. (London)* A **150** 9